

ADVANTAGES

- Ready to use and CE/IVD marked
- Simple preparative phase
- Maximum specificity
- Recovery of analytes ranges from 90% to 100%
- Stable for 3 years after production

*Accurate, Reproducible
and Quick Results*



CONTACT US

Eureka S.r.l. Lab Division
Via Enrico Fermi, 25
60033 Chiaravalle (AN) Italy

Tel: +39 071 745 0790

Fax: +39 071 749 6579

e-mail:

info@eurekaone.com



Website



Follow us on



OCCUPATIONAL TOXICOLOGY

September 2022



HPLC



IFU

URINE

- 1-Hydroxypyrene
- Acetone
- Free 2,5 Hexanedione
- Total 2,5 Hexanedione
- Hippuric Acid/o-m-p- Methylhippuric Acids
- Methanol
- Methyl-Ethyl-Keton (MEK)
- o-Cresol and Phenol
- Phenylglyoxilic Acid/Mandelic Acid
- Phenylglyoxilic/Mandelic/Hippuric/ o-m-p-Methylhippuric Acids
- S-phenylmercapturic Acid
- Trans, trans-Muconic Acid (ttMA)

WHOLE BLOOD

- Free Eritrocitary Protoporphyrin IX and ZnPP

GC, GC/MS



IFU

URINE

- Acetates
- Butyl Acetate
- Butyl Alcohol
- Carbonium Tetrachloride
- Diethylketone
- Ethyl Acetate
- Formic Acid
- Isobutyl Acetate
- Isobutyl Alcohol
- Isopropyl Alcohol
- Methyl Acetate
- Methylene Chloride (Dichloromethane)
- Methyl-Isobutylketone
- Methyltrichloroacetate
- N-Methylacetamide and N-Methylformamide
- Propyl Alcohol
- Tetrachloroethylene
- Trans-1,2-Cyclohexanediol
- Trichloroacetic Acid
- Trichloroethane
- Trichloroethanol
- Trichloroethylene
- Volatile Organic Compounds (VOC)

WHOLE BLOOD

- Volatile Organic Compounds (VOC)


LC-MS/MS



IFU

URINE

- Muconic Acid and S-Phenylmercapturic Acid



Control in urine - Level 2
S-Phenylmercapturic Acid (transition 238 – 109) 80 ng/ml (upper panel)
Deuterated Internal Standard (lower panel)